

Sleep Disorders and their Consequences on the Military Personnel: A Narrative Review

Sadegh Zarei ¹, Solmaz Sadat Hosseini Zijoud ² *

¹ Department of Clinical Biochemistry, School of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

² Department of Health Education and Health Promotion, School of Public Health, Iran University of Medical Sciences, Tehran, Iran

* **Corresponding Author:** Solmaz Sadat Hosseini Zijoud, Department of Health Education and Health Promotion, School of Public Health, Iran University of Medical Sciences, Tehran, Iran. Email: Hosseini1920@gmail.com

Received October 14, 2019; Accepted December 07, 2019; Online Published January 01, 2020

Abstract

Sleep is an integral part of the human life, and its disorders, depending on cognitive or emotional behaviors, cause a series of changes in the physical and mental activity of human beings. Therefore, as quality of sleep decreases, quality of life, performance and work efficiency will decrease as well. Military societies are in dire need of high quality sleep because of their duty of securing and protecting countries. For many reasons, including stressful and dangerous martial situations, sleep disorders are seen in both active and retired military personnel. The aim of the present study was to review the sleep disorders and their consequences on the military staff.

Keywords: Sleep Disorders, Sleep Quality, Military Personnel

Introduction

Sleep is a physiological and dynamic activity which reduces the awareness of the self and the environment. In the past, sleep was considered as a process where brain activity is reduced or completely eliminated.¹ During sleep, the body is physiologically active and a variety of dynamic fluctuations in the central nervous system and hemodynamic changes occur in the respiratory-ventilation and metabolic systems.² Sleep represents one-third of every person's life and has a tremendous impact on how people live and function throughout their lives. In fact, it is as important as the air we breathe and the food we eat.³ It is believed that good sleep strengthens health. Previous studies have shown that sleep disorders are common and associated with complications and mortality.⁴ Thus, sleep disorders in recent years have been considered as a branch of medicine, and their role in a number of seemingly unrelated clinical problems have been revealed.¹ In fact, sleep plays an important role in the rehabilitation and the focus of the individual's senses, and paying attention to its quality and quantity is considered to be extremely important.² Sleep disorders are also common in the military. In military societies, sleep disorders can lead to an increase in the Body Mass Index (BMI), non-compliance with sports standards, poor nutritional and health status, and most importantly, reduced military efficiency in the face of war, and other risks.⁵ These issues actually point out the importance of recognizing sleep disorders throughout every society as well as the military community. Therefore, in this review, sleep disorders and their consequences on the

military personnel have been studied.

Sleep Disorders

In today's stressful and competitive society, sleep deprivation or sleep disorders have gradually become a common social problem.⁶ Studies have shown that diseases such as anxiety and insomnia can cause severe injury and even death if left untreated.⁷

In 2014, the third edition of the International Classification of Sleep Disorders was published, which was comprised of six major groups: 1- Insomnia, 2- Sleep apnea syndrome (obstructive sleep apnea) 3- Centralized sleep and wake disorders 4- Circadian sleep-wake disorders 5- Parasomnia 6- Sleep-related movement disorders. This grouping is based on the main symptoms. The treatment of sleep disorders includes dietary changes, sleep hygiene, medication, and, in the case of sleep-dependent respiratory disorders, the use ventilators with positive and non-invasive pressure.⁸

However, sleep disorders (more than 90 types of disorders), each of which have a clear diagnostic criterion, can be divided into three major groups: 1- Nighttime insomnia (difficulty falling asleep, continuing to sleep or sleep without rest) 2- Daily drowsiness caused by nighttime insomnia 3- Sleep disturbing behaviors such as obstructive sleep apnea, parasomnia etc.⁹

Consequences of Insomnia in Military Personnel

Insomnia (dissatisfaction with the quality or quantity of sleep with complaints of falling asleep or continuing to sleep)

is one of the most common types of sleep disorders and a risk factor for a wide range of physiological and psychological disorders including depression, drug, antidepressants, etc. with obvious clinical anxiety and social, functional, and occupational problems.¹⁰ Cases such as old age, female gender (women twice as much as men) and poor socioeconomic status are the most common causes of insomnia, although their causes are not well-known.^{11,12} According to another study, about a third of the people in the West experience a number of insomnia symptoms each year.¹³ Depression is the most common cause of chronic insomnia.¹⁴ Of course, insomnia alone is not a diagnosis, but a symptom of a sleep disorder that has its own specific treatment.⁹

A person's vulnerability to sleep deprivation at various times, is different. Sleep deprivation can lead to many complications, such as increased risk of death, daily drowsiness, decreased consciousness, and hypothermia. Sleep deprivation is associated with increase in sympathetic activity, reducing cognitive functions such as attention and information processing, decision-making, and response time. The risk of coronary heart disease is higher in people with less than 6 hours of sleep than in people with a sleep period of more than 6 hours. Sleep deprivation also increases the hunger and appetite for salty foods and increases the risk of obesity in the military.⁵ Actually, insomnia is associated with decreased physical, mental, and cognitive impairment.^{14,15}

Bramoweth and Germain noted that insomnia is a common disorder in the military, especially those stationed in support of combat efforts. Situational stressors such as exposure to war, mild Traumatic Brain Injury (mTBI), irregular sleep and wake schedules, and delayed home return are all contributing to insomnia. However, insomnia is a risk factor for psychological stress reactions such as Post Traumatic Stress Disorder (PTSD), depression, suicide, and even the intent to kill other military or civilian people. Military people with mTBI are significantly affected by insomnia and have the majority of sleep disorders, which can impede their recovery, rehabilitation and military performance.¹⁶ Other similar studies suggest that insomnia affects 10-22 percent of civilians, while it is highly prevalent in active military personnel¹⁷ as well as retired veterans.¹⁸ Recent studies have shown a higher prevalence of insomnia among American men and women and Western allies who participated in the Operation Iraqi Freedom, compared to the civilian samples, 24-54%.¹⁸ In both military and civilian populations, insomnia is often associated with mental and physical health disorders such as PTSD, depression, brain injury and chronic pain.¹⁸ Many other studies have suggested that war PTSD has a major role in insomnia.¹⁹⁻²¹ Studies in Iran on chemical warfare veterans and their spouses showed insomnia and

poor sleep quality in these people due to PTSD in war veterans and excessive fatigue and confusion in their spouses. Studies also found that poor sleep quality in these people led to their severe depression and high consumption of antidepressants.²²⁻²⁴

Consequences of Respiratory Sleep Apnea Syndrome on Military Personnel

Obstructive Sleep Apnea Syndrome (OSAS) is a relatively common sleep-related disorder with a prevalence of 2-4%, characterized by repeated periods of partial or complete upper airway obstruction during sleep. It occurs in following arterial oxygen deprivation, cerebral arousal, and respiratory effort to end respiratory failure. The main clinical findings include daytime sleepiness and long night snoring.²⁵⁻²⁸ However, patients may also have other complaints such as morning headaches, insomnia, or nightmares. The diagnosis is nocturnal polysomnography (PSG) and is the treatment of choice in adults with persistent positive airway pressure.^{25,28,29} Lack of timely diagnosis can lead to a range of serious complications, including hypertension and systemic hypertension, cardiovascular disease, cognitive impairment, and psychiatric symptoms such as depression, irritability, and anxiety.³⁰⁻³³ Due to the nature of the disease that respiratory events occur during sleep, the diagnosis is usually delayed.³⁰ The US Department of Defense medical information has reported a significant increase in the diagnosis of insomnia and Obstructive Sleep Apnea (OSA) in all military services. From 2001 to 2009, OSA detection rates increased 8.5-fold due to the nature of the disease.³⁰ The US Department of Defense medical information has reported a significant increase in the diagnosis of insomnia and Obstructive Sleep Apnea (OSA) in all military services. From 2001 to 2009, OSA detection rates increased 5.8-fold.²⁰ Sleep disorders are most likely the result of increased insomnia.^{34,35} However, previous studies using PSG on the military personnel showed that OSA appears primary.³⁶⁻³⁸ Primary insomnia and OSA are conditions which are initially called "insomnia complex".³⁹ So sleep apnea may be exacerbated by the harmful effects of insomnia and lack of sleep, which makes military personnel in need of care. Failure to fully address all aspects of sleep disorders can potentially jeopardize the ability of military personnel to perform their duties or fully respond to treatment.²⁰ Current studies have linked experiences or war zones to the development of physical and psychological problems, especially OSA, including TBI-related illnesses, PTSD disorder, depression etc.^{18,40-42} Mysliwiec et al, have shown that 58.1% of the military personnel under study had one or more symptoms of sleep disorders and that OSA patients were twice as likely as those with PTSD and 1.5 times more parasomnia likely to have physical and psychological pain.⁴³ Collen et al. reported

that 97.4% of TBI soldiers complained of OSA as a primary diagnosis. A more comprehensive assessment of sleep disorders and related diagnoses may have important implications for the health and well-being of the military personnel.³⁵

Consequences of Sleep-related Movement Disorders on Military Personnel

Restless Legs Syndrome (RLS), Periodic Limb Movement Disorder (PLMD), Sleep-related Leg Cramps, Sleep-related Cramping Teeth, and Sleep Rhythmic Movement Disorder (RMD) are classified as Sleep-Related Movement Disorders (SRMD) (44). These clinical conditions are characterized by relatively simple, usually stereotyped, movements that can disrupt sleep and cause insomnia, poor sleep quality, fatigue and excessive daytime sleepiness. Physiological movements such as hypnotic jumps may occur during sleep. In some patients, the frequency and intensity of these movements may increase and cause sleep disorders.⁴⁴ The RLS is a sleep disorder characterized by four pathognomonic clinical features: (a) Involuntary movement of the legs, which is an unpleasant and inappropriate movement in the legs. (b) Unpleasant feeling that gets worse when you move or insist on moving. (c) Unpleasant feeling that is reduced by a slight shift or insistence on movement. (d) Unpleasant feeling in insisting on moving that is worse at night than during the day. These symptoms represent four basic criteria for detecting the presence of RLS.⁴⁵ Several epidemiological studies have evaluated the occurrence of RLS in the general population. These studies have observed a large variation in prevalence between 0.1% and 15%.⁴⁶ Higher prevalence has been reported in certain conditions (from high to low) respectively, in renal failure,⁴⁷ pregnancy,⁴⁸ and neuropathy.⁴⁹ Also, RLS is associated with depression.⁵⁰ As these conditions are common in military and veterans with PTSD,⁵¹ it is important to study the symptoms of RLS in these individuals. One study revealed that the prevalence of RLS was 1.16% in the elderly and 15.15% in the Chinese veterans. Also, the prevalence of RLS in the Chinese veterans was higher than their foreign counterparts. The prevalence of RLS in the elderly increases significantly with age, but the severity of RLS symptoms does not correlate with advancing age. Since the rate of diagnosis and treatment for RLS adults is extremely low in China and other countries, more attention should be paid to RLS knowledge.⁵² In a study by Rezaeitalab et al.⁵³ on thirty-one Iranian military veterans with PTSD it was revealed that all veterans had OSA while the prevalence of RLS was 22.5%. The high prevalence of RLS in these individuals suggests that PTSD in veterans may be associated with RLS. Insomnia, nightmares, sleep-related movement disorders and daytime sleepiness are often reported by PTSD veterans.⁵⁴ Studies using PSG have shown that the high

coefficient of PLMD is a sleep-related disorder in veterans with PTSD. Since PLMS is the most important target finding for RLS, it can be expected that RLS is high in these patients.⁵⁴⁻⁵⁶

Consequences of Parasomnia on Military Personnel

Parasomnia is a group of sleep disorders that includes unwanted events or experiences that occur when you start sleeping, sleeping, or waking up. Parasomnia may include abnormal movements, behaviors, emotions, perceptions, or dreams. Although the behaviors may be complicated and seemingly purposeful to others, you sleep during the event and often have no recollection of what has happened. If you have parasomnia, it is difficult to sleep at night.⁵⁷ Nightmares and disturbing night behaviors that occur after traumatic experiences have had various clinical features that overlap with other diagnoses of parasomnia. Nightmares with frequent occurrence of long, very lifeless dreams that cause significant distress or clinical disturbance in the social, occupational, or other function are prominent features of Parasomnia.⁴⁴ Occasional nightmares are part of a normal response to life stressors, with 85% of adults reporting at least one nightmare per year.⁵⁸ Participants from World War II generation had a significantly higher incidence of nightmares (7.2% in men and 7% in women). However, in normal populations this rate was 3.5% in men and 4.8% in women.⁵⁹ Nightmares are more common among patients with sleep disorders, depression, anxiety, and PTSD.⁶⁰⁻⁶² In addition, nightmares are associated with an increased risk of suicide in the military and civilian population.^{63,64} As sleep disorders are frequently reported in the US military personnel, this population may have a high incidence of nightmare disorder (NDO).²⁰ However, most research on the evaluation of sleep disorders in active duty military personnel has focused on insomnia, OSA, and PTSD.^{20,65} Following traumatic and stressful experiences, sleep disorders, including Trauma-Related Nightmares (TRN), have been reported.⁶⁵ The experience of nightmares is usually in the form of severe mental stress associated with periods of sleep disorder or deprivation. A limited number of laboratory tested cases have been reported with Eye Movement Disorder (RBD) and PTSD. Although not unique to PTSD, TRN has been recognized as a "hallmark" of the disorder and has a rate of up to 90%.^{66,67} The TSD including provocative traumatic experiences and the clinical features of trauma-related nightmares and disturbing psychotic nocturnal behaviors, is recognized as a specific parasomnia.⁶⁸ It should be noted that the physiology of PTSD traumatic nightmares is different compared to natural dreams. Traumatic nightmares are largely seen during sleep and the disrupt Rapid Eye Movement (REM) and are often associated with movement activity.⁴³ The TSD meets the basic criteria for a parasomnia

defined in the International Classification of Sleep Disorders because it includes abnormal sleep, sleep-related movements, behaviors, and autonomic nervous system activity that are not explained by another clinical disorder.⁴⁴ A study by Mysliwiec et al. reported a very low incidence of parasomnia among military personnel, with only 6 out of 725 military personnel suffering from various symptoms of parasomnia such as nightmares and sexual dysfunction. Therefore, there is a need for more studies on this group in the military community compared to civilians.³⁸

Consequences of Narcolepsy (Sleep Attack) on Military Personnel

Chronic narcolepsy disorder is seen primarily as a tendency to fall asleep and most cataplexy (sudden loss of coordination and muscle tone during waking). Cataplexy is generally mild and is caused by severe emotions. These emotions are usually positive (e.g. laughter), but different emotions have been reported. Patients with narcolepsy also experience disturbed night sleep, sleep paralysis, hallucinations in sleep and wakefulness.^{44,69} The clinical presentation of narcolepsy varies with age and its exact cause is not fully understood. Its causes can be environmental (stress, trauma, exposure to toxins and infection), genetic or autoimmune.⁷⁰⁻⁷² People with narcolepsy may tend to be overweight. This is not fully understood, but may be due to lower levels of physical activity, diet, metabolism, medication use, or a combination of these. Narcolepsy is also associated with an increased risk of depression. This may be because the symptoms of narcolepsy can be stressful.⁷³ Baseline Sleep Characteristics were explored for 71 US military service members with mTBI and indicated that 32% were suffering from restless legs syndrome and 6% were suffering from catalepsy. Older participants and those who were more aggressive, anxious or depressed had higher risks of OSA, catalepsy, etc.⁷⁴ Another study on 200 veterans and 20 former prisoners of war found that although exogenous factors (trauma) play an important role in accelerating PTSD and narcolepsy, but there is a genetic marker for narcolepsy in both groups. It has also been shown in this study that with the onset of PTSD, narcolepsy can be resistant and progressive.⁷⁵ Vergun reported that two most common sleep disorders in the military are insomnia and obstructive sleep apnea, whereas narcolepsy is not a common sleep disorder among the military personnel.⁷⁶

There has been no study on the other unexplained groups of sleep disorders in the military community. Of course, given the evidence of these unreported groups of disorders, one can guess that there is a high outbreak in active military and retired veterans following stressful events, especially war and stressful military situations. Studies on sleep disorders in the military community in Iran are also extremely limited, so further studies are needed to address the problem of military

disorders in order to provide solutions or treatments to increase the efficiency of the military personnel.

Conclusion

Sleep is extremely important for the operational readiness of the military personnel both operational and support, but is often overlooked during operational planning. Sleep disorders have serious consequences for the psychological, physical, and practical preparation both now and in the future. This means promoting healthy sleep and preventing sleep disorders may increase the readiness and resilience of the forces. The planning and management of sleep gives military units and their commanders the benefits of maintaining alertness, the medium-term tactical advantage of reduced sleep sensitivity and behavioral health disorders, and a long-term strategic advantage by increasing the efficiency of consciousness. Although the majority of publications have mentioned that seven to eight hours of sleep is considered to be standard for both the military and normal people, there is little guidance on how to achieve quality sleep.

Acknowledgments

None.

Authors' Contribution

All authors pass the four criteria for authorship contribution based on the International Committee of Medical Journal Editors (ICMJE) recommendations.

Conflict of Interests

The authors declared no potential conflict of interests with respect to the research, authorship, and/or publication of this article.

Funding/Support

The authors received no financial funding or support for the research.

References

1. Lee-Chiong. Sleep medicine essentials: John Wiley & Sons; 2011. p1-34.
2. Jaliloghadr S. Sleep and its disorders in Pediatrics. Qazvin: Qazvin Univ Med Sci; 2012.[Book]
3. Chapman DP, Wheaton AG, Perry GS, Sturgis SL, Strine TW, Croft JB. Household demographics and perceived insufficient sleep among US adults. *Journal of community health*. 2012;37(2):344-9. doi: 10.1007/s10900-011-9451-x
4. He J, Kryger MH, Zorick FJ, Conway W, Roth T. Mortality and apnea index in obstructive sleep apnea: experience in 385 male patients. *Chest*. 1988;94(1):9-14. doi.org/10.1378/chest.94.1.9
5. Lentino CV, Purvis DL, Murphy KJ, Deuster PA. Sleep as a component of the performance triad: the importance of sleep in a military population. *US Army Medical Department Journal*. 2013 Oct 1.
6. Trevor AJ, WL W. Sedative-hypnotic drugs. In: Katzung BG, Masters SB, AJ T, editors. *Basic & clinical pharmacology*. New York: The McGraw-Hill Companies; 2012. p. 482-95.[Book]

7. Vogliardi S, Favretto D, Tucci M, Stocchero G, Ferrara SD. Simultaneous LC-HRMS determination of 28 benzodiazepines and metabolites in hair. *Analytical and bioanalytical chemistry*. 2011;400(1):51-67. doi: [10.1007/s00216-011-4742-8](https://doi.org/10.1007/s00216-011-4742-8)
8. Onka K, Susta M. Past and present in drug treatment of sleep disorders. *Neuroendocrinology Letters*. 2014;35(3).
9. Pagel JF, Parnes BL. Medications for the treatment of sleep disorders: an overview. Primary care companion to the *Journal of clinical psychiatry*. 2001;3(3):118. doi: [10.4088/pcc.v03n0303](https://doi.org/10.4088/pcc.v03n0303)
10. Castillo R, Carlat D, Millon T, Millon C, Meagher S, Grossman S, et al. *Diagnostic and statistical manual of mental disorders*. American Psychiatric Association Press, Washington, DC. 2007.
11. Lichstein KL, Taylor DJ, McCrae CS, ME P. Insomnia: Epidemiology and risk factors. In: Kryger MH, Roth T, WC D, editors. *Principles and practice of sleep medicine*. 6th ed. Philadelphia: Elsevier Inc; 2017. p. 761-8. [Book]
12. Jausset I, Dauvilliers Y, Ancelin ML, Dartigues JF, Tavernier B, Touchon J, Ritchie K, Besset A. Insomnia symptoms in older adults: associated factors and gender differences. *The American Journal of Geriatric Psychiatry*. 2011;19(1):88-97. doi: [10.1097/JGP.0b013e3181e049b6](https://doi.org/10.1097/JGP.0b013e3181e049b6)
13. Ellis JG, Perlis ML, Neale LF, Espie CA, Bastien CH. The natural history of insomnia: focus on prevalence and incidence of acute insomnia. *Journal of psychiatric research*. 2012;46(10):1278-85. doi: [10.1016/j.jpsychires.2012.07.001](https://doi.org/10.1016/j.jpsychires.2012.07.001)
14. Roth T, Coulouvrat C, Hajak G, Lakoma MD, Sampson NA, Shahly V, Shillington AC, Stephenson JJ, Walsh JK, Kessler RC. Prevalence and perceived health associated with insomnia based on DSM-IV-TR; international statistical classification of diseases and related health problems, tenth revision; and research diagnostic criteria/international classification of sleep disorders, criteria: results from the America insomnia survey. *Biological psychiatry*. 2011;69(6):592-600. doi: [10.1016/j.biopsych.2010.10.023](https://doi.org/10.1016/j.biopsych.2010.10.023)
15. Pace-Schott EF, Spencer RM. Age-related changes in the cognitive function of sleep. In *Progress in brain research* 2011 Jan 1 (Vol. 191, pp. 75-89). Elsevier. doi: [10.1016/B978-0-444-53752-2.00012-6](https://doi.org/10.1016/B978-0-444-53752-2.00012-6)
16. Bramoweth AD, Germain A. Deployment-related insomnia in military personnel and veterans. *Current psychiatry reports*. 2013;15(10):401. doi: [10.1007/s11920-013-0401-4](https://doi.org/10.1007/s11920-013-0401-4)
17. Cozza SJ, Benedek DM, Bradley JC, Grieger TA, Nam TS, Waldrep DA. Topics specific to the psychiatric treatment of military personnel. *Iraq war clinician guide*. 2004;2:4-20.
18. Hoge CW, McGurk D, Thomas JL, Cox AL, Engel CC, Castro CA. Mild traumatic brain injury in US soldiers returning from Iraq. *New England journal of medicine*. 2008;358(5):453-63. doi: [10.1056/NEJMoa072972](https://doi.org/10.1056/NEJMoa072972)
19. McLay RN, Klam WP, Volkert SL. Insomnia is the most commonly reported symptom and predicts other symptoms of post-traumatic stress disorder in US service members returning from military deployments. *Military medicine*. 2010;175(10):759-62. doi: [10.7205/milmed-d-10-00193](https://doi.org/10.7205/milmed-d-10-00193)
20. Mysliwiec V, Gill J, Lee H, Baxter T, Pierce R, Barr TL, Krakow B, Roth BJ. Sleep disorders in US military personnel: a high rate of comorbid insomnia and obstructive sleep apnea. *Chest*. 2013;144(2):549-57. doi: [10.1378/chest.13-0088](https://doi.org/10.1378/chest.13-0088)
21. Wright KM, Britt TW, Bliese PD, Adler AB, Picchioni D, Moore D. Insomnia as predictor versus outcome of PTSD and depression among Iraq combat veterans. *Journal of clinical psychology*. 2011;67(12):1240-58. doi: [10.1002/jclp.20845](https://doi.org/10.1002/jclp.20845)
22. Tavalalaie SA, Assari SH, Najafi M, Habibi M, Ghanei M. Study of sleep quality in chemical-warfare-agents exposed veterans. *Journal Mil Med*. 2005;6(4):241-8.
23. Hojjati H, Ebadi A, Akhondzadeh G, ZARE K, Sirati M, Heravi M, NOHI E. Sleep quality in spouses of war veterans with post-traumatic stress: A qualitative study. 2017; 4(1): 1-9. doi: [10.29252/mcs.4.1.1](https://doi.org/10.29252/mcs.4.1.1)
24. SS H. The Relationship and Predicting Role of Sleep Quality and Sexual Self-esteem in Secondary Post-traumatic Stress Disorder among Wives of Veterans. *Military Caring Sciences Journal*. 2019;6(3):193-203. doi: [10.29252/mcs.6.3.4](https://doi.org/10.29252/mcs.6.3.4)
25. Azagra-Calero E, Espinar-Escalona E, Barrera-Mora JM, Llamas-Carreras JM, Solano-Reina E. Obstructive sleep apnea syndrome (OSAS). Review of the literature. *Medicina oral, patologia oral y cirugia bucal*. 2012;17(6):e925. doi: [10.4317/medoral.17706](https://doi.org/10.4317/medoral.17706)
26. Drager LF, Polotsky VY, Lorenzi-Filho G. Obstructive sleep apnea: an emerging risk factor for atherosclerosis. *Chest*. 2011;140(2):534-42. doi: [10.1378/chest.10-2223](https://doi.org/10.1378/chest.10-2223)
27. Rezaeitalab F, Foroughipour M, Sasannejad P, Moharrari F, Farzadfar MT, Mousavi Mirzaei SM. The association of headaches with obstructive sleep apnea syndrome diagnose at the sleep laboratory of Ebn-e-Sina hospital in Mashhad. *medical journal of mashhad university of medical sciences*. 2014;56(6):323-9.
28. BaHammam AS, Al-Shimemeri SA, Salama RI, Sharif MM. Clinical and polysomnographic characteristics and response to continuous positive airway pressure therapy in obstructive sleep apnea patients with nightmares. *Sleep medicine*. 2013;14(2):149-54. doi: [10.1016/j.sleep.2012.07.007](https://doi.org/10.1016/j.sleep.2012.07.007)
29. Kryger MH, Roth T, Dement WC. *Principles and practice of sleep medicine*. Elsevier; 2017. [Book]
30. McNicholas WT, Bonsignore MR, Management Committee of EU Cost Action B26. Sleep apnoea as an independent risk factor for cardiovascular disease: current evidence, basic mechanisms and research priorities. *European Respiratory Journal*. 2007;29(1):156-78. doi: [10.1183/09031936.00027406](https://doi.org/10.1183/09031936.00027406)
31. Lurie A. Cardiovascular disorders associated with obstructive sleep apnea. In *Obstructive Sleep Apnea in Adults*. *Advances in Cardiology*. 2011; 46: 197-266. doi: [10.1159/000325110](https://doi.org/10.1159/000325110).
32. Kielb SA, Ancoli-Israel S, Rebok GW, Spira AP. Cognition in obstructive sleep apnea-hypopnea syndrome (OSAS): current clinical knowledge and the impact of treatment. *Neuromolecular medicine*. 2012;14(3):180-93. doi: [10.1007/s12017-012-8182-1](https://doi.org/10.1007/s12017-012-8182-1)
33. Rezaeitalab F, Moharrari F, Saberi S, Asadpour H, Rezaeitalab F. The correlation of anxiety and depression with obstructive sleep apnea syndrome. *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences*. 2014; 19(3):205.
34. Luxton DD, Greenburg D, Ryan J, Niven A, Wheeler G, Mysliwiec V. Prevalence and impact of short sleep duration in redeployed OIF soldiers. *Sleep*. 2011;34(9):1189-95. doi: [10.5665/SLEEP.1236](https://doi.org/10.5665/SLEEP.1236)
35. Collen J, Orr N, Lettieri CJ, Carter K, Holley AB. Sleep disturbances among soldiers with combat-related traumatic brain injury. *Chest*. 2012;142(3):622-30. doi: [10.1378/chest.11-1603](https://doi.org/10.1378/chest.11-1603)
36. Capaldi VF, Guerrero ML, Killgore WD. Sleep disruptions among returning combat veterans from Iraq and Afghanistan. *Military medicine*. 2011;176(8):879-88. doi: [10.7205/milmed-d-10-00440](https://doi.org/10.7205/milmed-d-10-00440)
37. Lettieri CJ, Eliasson AH, Andrada T, Khramtsov A, Raphaelson M, Kristo DA. Obstructive sleep apnea syndrome: are we missing an at-risk population?. *Journal of Clinical Sleep Medicine*. 2005;1(04):381-5. doi: [10.1378/chest.128.4_MeetingAbstracts.1335-b](https://doi.org/10.1378/chest.128.4_MeetingAbstracts.1335-b)
38. Mysliwiec V, McGraw L, Pierce R, Smith P, Trapp B, Roth BJ. Sleep disorders and associated medical comorbidities in active duty military personnel. *Sleep*. 2013;36(2):167-74. doi: [10.5665/sleep.2364](https://doi.org/10.5665/sleep.2364)
39. Krakow B, Melendrez D, Pedersen B, Johnston L, Hollifield M, Germain A, Koss M, Warner TD, Schrader R. Complex insomnia: insomnia and sleep-disordered breathing in a consecutive series of crime victims with nightmares and PTSD. *Biological psychiatry*. 2001;49(11):948-53. doi: [10.1016/s0006-3223\(00\)01087-8](https://doi.org/10.1016/s0006-3223(00)01087-8)
40. Lew HL, Pogoda TK, Hsu PT, Cohen S, Amick MM, Baker E, Meterko M, Vanderploeg RD. Impact of the "polytrauma clinical triad" on sleep disturbance in a department of veterans affairs outpatient rehabilitation setting. *American Journal of Physical Medicine & Rehabilitation*. 2010;89(6):437-45. doi: [10.1097/phm.0b013e3181ddd301](https://doi.org/10.1097/phm.0b013e3181ddd301)
41. Seal KH, Bertenthal D, Miner CR, Sen S, Marmar C. Bringing the war back home: Mental health disorders among 103 788 US veterans returning from Iraq and Afghanistan seen at Department of Veterans Affairs Facilities. *Archives of internal medicine*. 2007;167(5):476-82. doi: [10.1001/archinte.167.5.476](https://doi.org/10.1001/archinte.167.5.476)
42. Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL. Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *New England Journal of Medicine*. 2004;351(1):13-22. doi: [10.1056/NEJMoa040603](https://doi.org/10.1056/NEJMoa040603)
43. Mysliwiec V, O'Reilly B, Polchinski J, Kwon HP, Germain A, Roth BJ. Trauma associated sleep disorder: a proposed parasomnia encompassing disruptive nocturnal behaviors, nightmares, and REM without atonia in trauma survivors. *Journal of Clinical Sleep Medicine*. 2014;10(10):1143-8. doi: [10.5664/jcs.m.4120](https://doi.org/10.5664/jcs.m.4120)
44. *International Classification of Sleep Disorders (ICSD-2): diagnostic and coding manual*. 2nd ed. Westchester: American Academy of Sleep Medicine 2006. [Book]

45. Allen RP, Picchietti D, Hening WA, Trenkwalder C, Walters AS, Montplaisi J. Restless legs syndrome: diagnostic criteria, special considerations, and epidemiology: a report from the restless legs syndrome diagnosis and epidemiology workshop at the National Institutes of Health. *Sleep medicine*. 2003;4(2):101-19. doi: [10.1016/s1389-9457\(03\)00010-8](https://doi.org/10.1016/s1389-9457(03)00010-8)
46. Merlino G, Valente M, Serafini A, Gigli GL. Restless legs syndrome: diagnosis, epidemiology, classification and consequences. *Neurol Sci*. 2007;28 Suppl 1:S37-46. doi: [10.1007/s10072-007-0736-x](https://doi.org/10.1007/s10072-007-0736-x)
47. Lin Z, Zhao C, Luo Q, Xia X, Yu X, Huang F. Prevalence of restless legs syndrome in chronic kidney disease: a systematic review and meta-analysis of observational studies. *Ren Fail*. 2016;38(9):1335-46. doi: [10.1080/0886022X.2016.1227564](https://doi.org/10.1080/0886022X.2016.1227564)
48. Chen SJ, Shi L, Bao YP, Sun YK, Lin X, Que JY, et al. Prevalence of restless legs syndrome during pregnancy: A systematic review and meta-analysis. *Sleep Med Rev*. 2018;40:43-54. doi: [10.1016/j.smrv.2017.10.003](https://doi.org/10.1016/j.smrv.2017.10.003)
49. Harashima SI, Nishimura A, Osugi T, Wang Y, Liu Y, Takayama H, Inagaki N. Restless legs syndrome in patients with type 2 diabetes: effectiveness of pramipexole therapy. *BMJ supportive & palliative care*. 2016;6(1):89-93. doi: [10.1136/bmjspcare-2014-000691](https://doi.org/10.1136/bmjspcare-2014-000691)
50. Koo BB, Blackwell T, Lee HB, Stone KL, Louis ED, Redline S, Osteoporotic Fractures in Men (MrOS) Study Group. Restless legs syndrome and depression: effect mediation by disturbed sleep and periodic limb movements. *The American Journal of Geriatric Psychiatry*. 2016;24(11):1105-16. doi: [10.1016/j.jagp.2016.04.003](https://doi.org/10.1016/j.jagp.2016.04.003)
51. Kibler JL, Tursich M, Ma M, Malcolm L, Greenberg R. Metabolic, autonomic and immune markers for cardiovascular disease in posttraumatic stress disorder. *World journal of cardiology*. 2014;6(6):455. doi: [10.4330/wjc.v6.i6.455](https://doi.org/10.4330/wjc.v6.i6.455)
52. Zhao L, Hu F. An epidemiologic study of restless leg syndrome among retired military elders in Shanxi Province. *Zhonghua yi xue za zhi*. 2015;95(5):382-5. doi: [10.3760/cma.j.issn.0376-2491.2015.05.016](https://doi.org/10.3760/cma.j.issn.0376-2491.2015.05.016)
53. Rezaeitalab F, Ardani AR, Ravanshad Y, Rezaeitalab F, Ardakani SA. Restless Leg Syndrome is Prevalent in Military Veterans with Post-Traumatic Stress Disorder and Sleep Disturbances. *J Trauma Treat*. 2018;7(413):2167-1222. doi: [10.4172/2167-1222.1000413](https://doi.org/10.4172/2167-1222.1000413)
54. Tipton PW, Wszolek ZK. Restless legs syndrome and nocturnal leg cramps: a review and guide to diagnosis and treatment. *Pol Arch Intern Med*. 2017;127(12):865-72. doi: [10.20452/pamw.4148](https://doi.org/10.20452/pamw.4148)
55. Germain A, Nielsen TA. Sleep pathophysiology in posttraumatic stress disorder and idiopathic nightmare sufferers. *Biological psychiatry*. 2003;54(10):1092-8. doi: [10.1016/s0006-3223\(03\)00071-4](https://doi.org/10.1016/s0006-3223(03)00071-4)
56. Ferri R, Koo BB, Picchietti DL, Fulda S. Periodic leg movements during sleep: phenotype, neurophysiology, and clinical significance. *Sleep medicine*. 2017;31:29-38. doi: [10.1016/j.sleep.2016.05.014](https://doi.org/10.1016/j.sleep.2016.05.014)
57. Parasomnias & Disruptive Sleep Disorders Cleveland: Cleveland Clinic; 2017 [Available from: <https://my.clevelandclinic.org/health/diseases/12133-parasomnias--disruptive-sleep-disorders>].
58. Levin R, Nielsen TA. Disturbed dreaming, posttraumatic stress disorder, and affect distress: a review and neurocognitive model. *Psychological bulletin*. 2007;133(3):482. doi: [10.1037/0033-2909.133.3.482](https://doi.org/10.1037/0033-2909.133.3.482)
59. Sandman N, Valli K, Kronholm E, Ollila HM, Revonsuo A, Laatikainen T, Paunio T. Nightmares: prevalence among the Finnish general adult population and war veterans during 1972-2007. *Sleep*. 2013;36(7):1041-50. doi: [10.5665/sleep.2806](https://doi.org/10.5665/sleep.2806)
60. Schredl M. Nightmare frequency and nightmare topics in a representative German sample. *European archives of psychiatry and clinical neuroscience*. 2010;260(8):565-70. doi: [10.1007/s00406-010-0112-3](https://doi.org/10.1007/s00406-010-0112-3)
61. Li SX, Lam SP, Yu MW, Zhang J, Wing YK. Nocturnal sleep disturbances as a predictor of suicide attempts among psychiatric outpatients: a clinical, epidemiologic, prospective study. *The Journal of clinical psychiatry*. 2010. doi: [10.4088/JCP.09m05661gry](https://doi.org/10.4088/JCP.09m05661gry)
62. Li S, Lam J, Zhang J, Yu M, Wing Y. Prevalence and correlates of nightmare disturbances in patients with sleep disorders other than REM-parasomnias. *Sleep Medicine*. 2013;14:e182-3. doi: [10.1016/j.sleep.2013.11.430](https://doi.org/10.1016/j.sleep.2013.11.430)
63. Lande RG. Sleep problems, posttraumatic stress, and mood disorders among active-duty service members. *The Journal of the American Osteopathic Association*. 2014;114(2):83-9. doi: [10.7556/jaoa.2014.021](https://doi.org/10.7556/jaoa.2014.021)
64. Nadorff MR, Nazem S, Fiske A. Insomnia symptoms, nightmares, and suicidal ideation in a college student sample. *Sleep*. 2011;34(1):93-8. doi: [10.1093/sleep/34.1.93](https://doi.org/10.1093/sleep/34.1.93)
65. Lettieri CJ, Williams SG, Collen JF. OSA syndrome and posttraumatic stress disorder: clinical outcomes and impact of positive airway pressure therapy. *Chest*. 2016;149(2):483-90. doi: [10.1378/chest.15-0693](https://doi.org/10.1378/chest.15-0693)
66. Ross RJ, Ball WA, Sullivan KA, Caroff SN. Sleep disturbance as the hallmark of posttraumatic stress disorder. *The American journal of psychiatry*. 1989. doi: [10.1176/ajp.146.6.697](https://doi.org/10.1176/ajp.146.6.697)
67. Writer BW, Meyer EG, Schillerstrom JE. Prazosin for military combat-related PTSD nightmares: a critical review. *The Journal of neuropsychiatry and clinical neurosciences*. 2014;26(1):24-33. doi: [10.1176/appi.neuropsych.13010006](https://doi.org/10.1176/appi.neuropsych.13010006)
68. Mysliwiec V, Brock MS, Creamer JL, O'Reilly BM, Germain A, Roth BJ. Trauma associated sleep disorder: a parasomnia induced by trauma. *Sleep medicine reviews*. 2018;37:94-104. doi: [10.1016/j.smrv.2017.01.004](https://doi.org/10.1016/j.smrv.2017.01.004)
69. Scammell TE. Narcolepsy. *New England Journal of Medicine*. 2015;373(27):2654-62. doi: [10.1056/NEJMra1500587](https://doi.org/10.1056/NEJMra1500587)
70. Nishino S, Ripley B, Overeem S, Lammers GJ, Mignot E. Hypocretin (orexin) deficiency in human narcolepsy. *The Lancet*. 2000;355(9197):39-40. doi: [10.1016/S0140-6736\(99\)05582-8](https://doi.org/10.1016/S0140-6736(99)05582-8)
71. Schneider L, Mignot E. Diagnosis and management of narcolepsy. *In Seminars in neurology* 2017 Aug (Vol. 37, No. 04, pp. 446-460). Thieme Medical Publishers. doi: [10.1055/s-0037-1605554](https://doi.org/10.1055/s-0037-1605554)
72. Chabas D, Taheri S, Renier C, Mignot E. The genetics of narcolepsy. *Annual review of genomics and human genetics*. 2003;4(1):459-83. doi: [10.1146/annurev.genom.4.070802.110432](https://doi.org/10.1146/annurev.genom.4.070802.110432)
73. Depression, Weight Gain, and Health Complications: the National Sleep Foundation; 2016 [Available from: <https://www.sleepfoundation.org/narcolepsy/what-narcolepsy/depression-weight-gain-and-health-complications/>].
74. Walker JM, James NT, Campbell H, Wilson SH, Churchill S, Weaver LK. Sleep assessments for a mild traumatic brain injury trial in a military population. *Undersea & hyperbaric medicine: journal of the Undersea and Hyperbaric Medical Society, Inc*. 2016;43(5):549-66.
75. Mellman TA, Ramsay RE, Fitzgerald SG. Divergence of PTSD and narcolepsy associated with military trauma. *Journal of anxiety disorders*. 1991;5(3):267-72. doi: [10.1016/0887-6185\(91\)90007-G](https://doi.org/10.1016/0887-6185(91)90007-G)
76. DAVID V. Army Researcher: Sleep, Readiness Go Hand in Hand: ARMY NEWS SERVICE; 2017 [Available from: <https://www.defense.gov/Explore/News/Article/Article/1127928/army-researcher-sleep-readiness-go-hand-in-hand/>].