

# Glymphatic and Immun Systems of the Brain

Rengin Kosif<sup>1</sup>, Ibrahim Kurtul<sup>2</sup>, Sinem Kara<sup>3\*</sup>, Umit Atasever<sup>3</sup>, H. Kubra Koc Topcuoglu<sup>3</sup>

<sup>1</sup> Kirikkale University, Faculty of Medicine, Department of Anatomy, Kirikkale, Turkey

<sup>2</sup> Bolu Abant Izzet Baysal University, Faculty of Medicine, Department of Anatomy, Bolu, Turkey

<sup>3</sup> Bolu Abant Izzet Baysal University, Faculty of Medicine, Department of Interdisciplinary Neuroscience, Bolu, Turkey

\* **Corresponding Authors:** Sinem Kara, Bolu Abant Izzet Baysal University, Faculty of Medicine, Department of Interdisciplinary Neuroscience, Bolu, Turkey. E-mail: [sinem.kara.2009@gmail.com](mailto:sinem.kara.2009@gmail.com)

Received February 11, 2021; Accepted May 29, 2021; Online Published June 5, 2022

## Abstract

Even though brain lymphatic system has several peculiarities including the construction of the endothelial cells of the lymph vessels, in common with the lymphatic system of the body in general, studies have documented eminent discrepancy. The unique lymphatic system of the brain is called "the glymphatic system" due to its lymphatic system-like function and glial-like fluid flow. The collaborative functions of this system, and CSF and interstitial fluid has been documented. A recent discovery of this system has highlighted the requirement to review the diagnosis and treatment procedures of several neuronal diseases. With this discovery, anatomic, physiopathologic, immunologic, and genetic natures of this system have attracted attention of scientists. This review has summarized the accumulating information on the lymphatic system of the brain, particularly focusing on the function of the glymphatic system in fetus and newborn, cranial tumors, neurodegenerative diseases, brain metabolism and its relationship with sleep, and the anatomical posture of the body. Yet, the cells responsible with the brain immune system and their relationship with the glymphatic system have been discussed. By summarizing the latest information on the relationship of this system with immunologic, molecular, physiologic, and anatomical aspects of several neuronal diseases, this paper will surely contribute to the diagnosis and treatment of them through shedding light on the researchers.

**Keywords:** Brain, Glymphatic System, Immunity, Neuronal Disease

## Introduction

Lymphatic system plays a critical role in homeostasis and function in peripheral tissue. Although the brain does not have a lymphatic vasculature in traditional thought, it has been understood that the brain contains a unique lymphatic formation among the somatic organs. No anatomical structures of the brain with very high metabolic functions and complexity have been determined before, facilitates the passage of fluid through the parenchyma and lymphatic clearance of extracellular substances.<sup>1</sup> Central Nerve System (CNS), previously thought to have no lymph system, has remained a mystery as a subject of research in the field of neuroscience for many years.<sup>2</sup>

The first definition of the lymphatic system on the brain goes back to Italian anatomist and illustrator Paolo Mascagni.<sup>3</sup> He clearly explained the Meningeal Lymph Vessels (MLV) and their functional roles more than 250 years ago, expressing them as "the privilege of the brain immunity". Another study carried out nearly 100 years later, indicated controversially detailed

explanation of the absence of a lymphatic circulatory system in the brain which misled many scientists for years.<sup>4</sup> Some reasons of this misconception have been cited as the presence of tight connections in Blood-Brain Barrier (BBB), lack of the conventional lymphatic drainage system, and data showing limited rejection of the foreign matter in CNS.<sup>5</sup>

With new data accumulated on the lymphatic system of the CNS, surprising findings have been obtained in recent years.<sup>5,7</sup> A study investigating T-cell gateways in the meninges, functional lymphatic vessels lining the subdural sinuses were discovered.<sup>8</sup>

This study revealed that MLVs are located specifically around the subdural sinuses at the base of the skull, and that the endothelial cells in these vessels show all the molecular features of their counterparts found in other lymphatic vessels and eventually attach to the deep cervical lymph nodes.<sup>9</sup> These lymphatic vessels take glial fluid and immune cells from the cerebrospinal fluid (CSF). Another study also has confirmed these

links.<sup>10</sup>

The lymphatic vessels of the CNS transport CSF throughout the perivascular and interstitial spaces. The CSF flow is then directed through venous perivascular spaces, and at the same time metabolic waste in the parenchyma through the CSF. This system is called “The Glymphatic System” because it functions like glial fluid flow and lymphatic system. It is indeed not a lymphoid tissue but is occasionally named as “lymphoid-like” tissue. These lymphatic vessels lie along the dura mater and advance throughout the superior sagittal sinus and peri-sinus space of the lateral subdural sinuses.<sup>6</sup>

Recent studies have proved that lymphatic vessels in the skull base of the rodents are responsible for draining the CSF and other waste materials out of the brain.<sup>11</sup> A study carried out on mice has shown that MLVs have turned into a complex network along the blood vessels in the base of the skull.<sup>12</sup> Yet, certain macromolecules infused into the lateral ventricle have been displayed to reach out the cervical lymph nodes using the foramen of the skull and perineural tracks.

Research have demonstrated, using magnetic resonance imaging, contrast agents injected into the interstitial fluid, absorbed by the basal meningeal vessels, and later drained out of the brain through the CSF.<sup>13,14</sup> A study has used Prox1–GFP to explain the function of the lymphatic vessels of the brain in mouse by marking the promotor region of the Prox1 gene, which has a key role in lymphatic development, with green fluorescent protein.<sup>12</sup> This study has displayed the unification pattern of the endothelial cells of the dorsal MLVs to be like closed zipper while that of the basal MLVs to be basically as a button, as is the case in the lymphatic capillary vessels of the peripheral organs.

In general, based on these anatomical and functional experiments, the delivery of macromolecules that cannot pass through the BBB in the brain to the peripheral lymphatic system and the drainage of macromolecules in the CSF are due to basal MLVs.<sup>12,14</sup>

### **Lymphatic Drainage System of the Brain in Fetus and Newborns**

Drainage of the interstitial fluid and CSF of the brain by the lymphatic system is essential in homeostasis. This drainage system may deteriorate with neurologic lacerations in both premature and full-term newborns. Brain damage due to acute hypoxia-ischemia in full-term newborns is an important death or injury cause.<sup>15</sup>

Various cellular and molecular pathologies such as neurovascular and BBB deformations, inflammation, damage in immune cells and lymphatic circulation, and deterioration in the brain restoration capacity, have been suggested in hypoxic-ischemic encephalopathy.<sup>16</sup> Astrocytes, important components of both BBB and the glymphatic system, are very essential mediator sources in the inflammation seen right after the cerebral ischemia.<sup>17</sup> An Magnetic Resonance Imaging (MRI) study has reported the meningeal lymphatic flow streaming in the sagittal sinus, in reverse direction with the venous flow.<sup>14</sup> Cerebral edema is the most apparent histological finding during the hypoxic ischemia. Animal studies have shown increased aquaporin-4 (AQP-4) expression in this finding after brain damage with different causes.<sup>18</sup> This increase has been associated with the development of the cerebral edema due to the increase in BBB permeability.<sup>19</sup> Therefore, acute brain damage ruins glymphatic function, in turn deteriorating itself due to accumulation of both normal and pathological metabolic wastes.<sup>15</sup>

The lymphatic drainage system of the brain has also been associated with several other pathologies such as white matter damage in newborns. Reactive astrogliosis is one of the regenerative mechanisms after the neuronal damage in the white matter of premature. It can change the nature and function of the glymphatic system. Waste metabolites produced during the oxidative stress and inflammation, representing main mechanism of the white matter development, are removed by MLVs, which is very vital in limiting neurological pathology in fetal development.<sup>15</sup>

### **Exclusive Immune System of the Brain**

Features that contribute to the opinion that CSN is immunologically privileged: mostly due to the presence of tight junctions in the BBB, the idea of the absence of a classical lymph drainage system, and limited rejection of foreign tissue in the CNS.<sup>5</sup> However, today the concept of immune privilege of the brain has been redefined. As mentioned above, functional lymph vessels have been described and antigen offering cells (APC) including microglia, macrophages, astrocytes, and Dendritic Cells (DCs) have been shown in CNS. It is known that these components are not isolated from T cells, being responsible in emptying CNS antigens locally into the cervical lymph nodes. The exclusive immune system

of CNS is not insufficient.<sup>5,20</sup> Either dysfunction or hyper function of the MLVs causes an autoimmune attack, resulting in destroyed meningeal immune activity with irregular ablations.<sup>7</sup> Hence, an autochthon production of the T cells and antibodies is present in the brain. Consequently, it is now accepted that the presence of the T cells in the brain is autoimmune based.<sup>9</sup>

Cytokinin production and migration of DCs to the lymph nodes have also been searched to reveal APC function of the CNS. In vitro studies have shown immunogenetic activity of DC.<sup>5,21</sup> Addition of certain agents into the medium such as type I interferon, synthetic Toll-like receptor 3 (TLR3) ligand, and polycytidylic acid has enhanced this activity. On the researches dealing with the inflammation that occurs in the vaccine area, granulocyte-macrophage colony stimulant factor (GM-CSF) 127, poly (I:C) 200 and other TLR agonists have been applied to the area to help increase the immunological response.<sup>5,22</sup> It is obvious, in the light of the information given above, that an isolated immunologic nature is present in the brain. Genetic risk factors are also important regulators of the lymphatic drainage system of the brain. In several of the cognitive Alzheimer Disease (AD) model, the diseases are caused by mutations in the macrophage gene expression.<sup>23,26</sup> APOE polymorphic alleles have been APOE major alleles considered as the primary genetic risk factor for AD.<sup>27,28</sup> Among the e2, e3, and e4, e4 has been suggested to be the strongest independent risk factor for the sporadic AD development.<sup>20</sup>

The new lymphatic concept of the brain has led to reconsider the autoimmune etiology of several neuro-inflammatory and neurodegenerative diseases including Multiple Sclerosis (MS) and AD.<sup>9</sup> Aging is considered as an important factor on the lymphatic drainage system of the brain. Fluid and metabolites produced by parenchyma of the brain may fast and effectively be eliminated by multi-lymphatic drainage system in the young. With increasing age, changes are observed in the cerebral vessel system. Atherosclerosis and decrease in the amplitude of pulsations lead to the fluid accumulation in the enlarged perivascular space, occlusion of the perivascular pathways along the basal membrane, and functional deterioration in the glymphatic drainage and CSF paths.<sup>29</sup>

Literature has reported cognitive inefficiency when MLVs are cut in young mice. MLVs have been documented

to have a close relation functionally with these kinds of pathologies including AD in neurodegenerative mouse models.<sup>7</sup> With degeneration of these vessels in aged mice, CSF drainage circulation has been deteriorated.<sup>6</sup> A similar study has shown eminent deteriorated dorsal MLVs and dilated and hyperplastic basal MLVs in the aged animals as compared to the young ones.<sup>12</sup> Another research has revealed a breakdown in the connections of the lymphatic endothelial cells because of the impairment in the lymphatic drainage, which is surely in relation with the lymphatic vessel dysfunction and degeneration due to aging.<sup>30</sup> Studies have striven to reveal age related changes. They have reported less zipper type connections and more button type ties between the lymphatic endothelial cells of the aged basal MLVs. These changes are possibly related to the decrease in the CSF drainage in aged mice. Another study using MS animal models has associated MS with MLV ablation, emphasizing the fact that MLVs need to be ameliorated to eliminate inflammation in the brain.<sup>6,12,30</sup>

Accumulation of toxic protein clusters is the common pathologic feature of neurodegenerative diseases including AD, Parkinson and Huntington diseases, and motor neuron disease. Amyloid- $\beta$  ( $A\beta$ ) accumulation is considered as the primary cause of AD. Recent findings have suggested that extracellular  $A\beta$  congregate is removed from the brain by several ways such as enzymatic degradation, phagocytosis by microglia, glycopytic clearance, and BBB and lymphatic ways. The last two ways have been shown to be particularly responsible for the clearing metabolic waste away from the brain.<sup>6,31</sup> Another report has revealed that ~40 % of the  $A\beta$  peptides and ~19 % of the pathologic tau protein has been removed from the brain through this way.<sup>32</sup> MLVs have also been reported to ease pathologies in relation with  $A\beta$  and tau protein in the brain through enhancing  $A\beta$  clearance,<sup>20,32</sup> therefore, suggesting a new therapeutic approach in neurodegenerative diseases.<sup>6</sup> After all, detail exploration of the brain lymphatic system may lead to new approaches to reveal connections between the brain parenchyma and peripheral tissues and related pathologies.<sup>6,33</sup>

There are several pathways between the brain and peripheral tissues for substance exchange such as BBB, choroid plexus, subarachnoid plexuses, and lymphoid system. Effectiveness of these ways decrease with the aging.<sup>34</sup> In a study, it has been suggested that there are

some physiological mechanisms that remove intracellular and extracellular pathological proteins from the brain, and play a critical role in the clearance of metabolic wastes and homeostasis. Displaying the vessels of the CNS with advanced technology may open a road to new approaches for the treatment of neuroinflammatory diseases.<sup>6,7</sup> Revealing the possible relation of the lymphatic system with the other clearance mechanisms is also essential. Likewise, determining the volume of the waste material cleared by the lymphatic system through the CSF may answer the question whether the brain lymphatic system plays an important role in either clearance of the waste materials or immune regulation. Basically, this question will bring about the possible role of the brain lymphatic system on the pathogenesis of neurodegenerative diseases, contributing to the development of the therapeutic agents.<sup>8,20</sup>

The glymphatic system is a perivascular space network supporting CSF flow and clearance of the waste materials.<sup>35</sup> Literature has indicated reciprocal transition of the CSF and brain interstitial fluid through the perivascular structures surrounding cerebral blood vessels.<sup>1</sup> This system is supported by AQP4 localized to the last vascular extensions of the astrocytes. Reports have documented three elements facilitating the CSF passage; the CSF flow path in periarterial space, the perivenous interstitial fluid clearance, and the astrocytic AQP4 transparenchymal channel.<sup>1,2</sup> The glymphatic system under the control of circadian rhythm is affected by the cardiovascular system and sleep, both being regulated by it. The space between the brain tissue and CSF compartments is broad enough to provide clearance by the basic diffusion. This is particularly the case for larger molecules such as proteins and peptides with lower diffusion coefficient.<sup>35</sup> A study has shown the relation of the glymphatic flow of the CSF from the cerebellomedullary cistern through the lymph nodes at day and night.<sup>35</sup> This data emphasizes the role of glymphatic system in the interstitial fluid elimination from the brain parenchyma during sleep.<sup>2</sup> This system is especially effective during sleep but whether sleep timing promotes it or not is unclear. It has clearly been reported that this system removes waste materials from the brain through increasing CSF activity.<sup>35,37</sup> In another study supporting these data, it has been revealed that the rodent brain was more effective at removing neuronal metabolism waste, especially during slow-wave sleep, compared to

the awake state.<sup>38</sup>

Brain energy metabolism may also play a role in regulating the function of the glymphatic system on the sleep-wakefulness cycle. Originally thought as a waste material, lactate may function, based on the hypothesis of the lactate cycle between astrocyte and neuron, as a supplementary fuel and a signal molecule in this metabolism. This idea is supported by the evidence revealing the presence of a lactate concentration gradient between the astrocyte and neuron.<sup>39,40</sup> Dynamic changes in the brain lactate level are regulated by the lymphatic clearance systems.<sup>36</sup> Sleep is known to have homeostatic function removing the neural metabolites.<sup>41</sup> It clears harmful metabolites from the brain such as  $A\beta$  that occurs during the wakefulness, and strengthen memory.<sup>42</sup> Consequently, enhancing the sleep quality may be helpful through lowering the development risk of the amyloid beta ( $A\beta$ ) pathology in AD protection.<sup>41</sup> Yet, acute insomnia ruins cognitive functions, and is usually the early result of most neurodegenerative diseases.<sup>42</sup>

### Effect of Body Posture on Lymphatic System Function

A research has examined the changes in CSF and interstitial fluid in rats at supine, facedown and lateral postures.<sup>43</sup> Among these positions, the lateral one representing sleep-resting in rats, have been shown to be the most effective posture in removing  $A\beta$  and other metabolites through glymphatic paths. Reversely, it is significantly low at the supine posture which is the ‘upright’ position at a conscious state in rats. The relationship between Intracranial Pressure (ICP) and body posture has been reported.<sup>20</sup> Another study has indicated ICP variations at different body postures in the idiopathic intracranial hypertension and mild hydrocephaly cases.<sup>44</sup>

### Intracranial Tumors

Mechanisms in MLVs are used to promote anti-tumor reaction. Several loaded antigen types are applied on DCs vaccine studies for brain tumors. EGFR<sup>vIII</sup> molecule functions are considered as an ideal tumor specific antigen for glioblastoma patients. It is actually expressed in nearly 30% of the patients. While vaccine is prepared, the EGFR<sup>vIII</sup> molecule on the T cell, a Chimeric Antigen Receptor (CAR), is marked. Detection of this molecule may be an indicator on therapy

options. CAR T cell is one of the immunologic indicators in lymphatic system. Decrease in the expression of that molecule in this cell provokes tumor recurrence. Accordingly, this complex system has been proved to clear off the brain tumors, and at long term can provide protection.<sup>45</sup> Another study has reported increases in cytokine production and chance of survival after the administration of the DC vaccine during the phase I and II trials of glioma patients.<sup>5</sup> In the light of this information, it can surely be pointed out that future immunotherapeutic studies will help us understand in detail the relationship between the metastatic tumors and lymphatic immune system of the brain.

The function of the brain lymphatic system is broadly affected by several pathological conditions. Changes of the compounds of constructive proteins in vascular diseases like lead to a decrease in vascular plasticity, deteriorating perivascular flow. Intracranial arterial narrowness not only restrain cerebral blood flow but also obstruct perivascular or paravascular pathways.<sup>46</sup> Traumatic brain damage, hemorrhagic or embolic paralyzes and diabetes destroy the glymphatic clearance of neurotoxic metabolites such as  $A\beta$ .<sup>20</sup> A study has determined irregularity in the expression polarity of the astrocytic AQP4 of the glymphatic system, right after the diffuse of ischemia in the multiple cerebral microinfarcts, leading to neurovascular-neurocognitive disorders.<sup>47</sup> Disorders seen in the lymphatic system drainage of the cerebrum due to either aging or different pathologies, lead to extracellular dyshomeostasis and antigen production. However, the dysfunction of this drainage system may lead to severe pathophysiological changes including neurodegenerative diseases, neuroinflammation and brain tumors, as reported by literature.<sup>20</sup>

Neurovascular disease and stroke are the main reasons of the geriatric disability. Secondary complications including cerebral aneurismal rupture, bleeding, hydrocephaly, and delayed cerebral ischemia related to micro thrombus, may also develop.<sup>48</sup> Lymphatic system drainage of the cerebrum can be disrupted by acute brain injuries, neurovascular diseases, and various stroke cases. This disruption may trigger secondary complications, ending up with more severe neuropathology. Functions of the cytokines produced by both macrophages and microglia in the neuroinflammation seen after ischemic stroke, have been documented.<sup>49,50</sup> There are two polarization phenotypes of the macrophages documented; one, classically activated

macrophages (M1 type), the other, alternatively activated macrophages (M2 type).<sup>51</sup> It has been reported that as ischemic infarcts develops, M1 phenotype produces proinflammatory cytokines, and M2 phenotype anti-inflammatory cytokines.<sup>49</sup> Besides, a broad variation is observed in the timing and contribution rates of the microglia/macrophage to the different phases of ischemic stroke. A study performed on mice has reported intense localized microglia very active at the first day of the acute ischemic stroke induced by the obstruction of the middle cerebral artery.<sup>52</sup> Blood originated macrophages have congregated thereby 3-7 days after the stroke. Spatially and temporally, the activation and polarization of microglia/macrophages that occur have been shown to coincide with dynamic changes in their phenotype during the development and progression of ischemic stroke. Perivascular spaces are essential regions for the initiation of neuroinflammation. Microglia may instantly become active as a reaction to hypoxia stimulation right after ischemic damage begins. Blood originated macrophages migrate to the paravascular spaces. Cytokines produced by the perivascular macrophages after the cerebral ischemia, have been thought to orientate infiltration of the inflammatory cells. Intravascular inflammation also activates intracerebral microglia after the subarachnoid hemorrhage. The activation and survival of the microglia induced by ischemic stimulus may be through the signals of Colony Stimulant Factor-1 (CSF-1) receptor.<sup>53</sup> Several research have examined the effects of the blockage of the lymphatic drainage in the cerebrum during the ischemic and hemorrhagic strokes.<sup>53,55</sup> Models of the Cervical Lymphatic Blockage (CLB) have been demonstrated to deteriorate the cerebral lymphatic drainage directly, leading to excessive fluid and waste material accumulation in the perivascular and paravascular spaces, ending up with the intracranial hypertension and brain edema.<sup>20</sup> Likewise, the effects of the blockage in the model of the ischemic stroke originating from the middle cerebral artery have been reported.<sup>54</sup> Brain edema and oxidative stress in the ischemic region has been documented in this literature. Additionally, severe neuronal damages have been observed. The blocking of drainage of the cervical lymphatic flow and middle cerebral artery has made the mentioned changes more noticeable. Another study has introduced a visualization procedure to measure interstitial fluid drainage of the brain dynamically at multiphoton microscope.<sup>56</sup> In the

stroke model induced by Rose Bengal which acts as a sensitizer against light through photo thrombosis, it has been shown to obstruct a single vessel segment, lowering the drainage of the interstitial fluid profoundly through deteriorating focal vascular perfusion, and not affecting the flow of other vessels.<sup>56</sup> Another report has documented the function of the glymphatic system in different stroke models through administrating contrast substance into the cerebellomedullary cistern, using MRI. The system has significantly been deteriorated in the acute phase of embolic ischemic stroke and after the subarachnoid hemorrhage.<sup>57</sup> These results have indicated the cerebral arterial pulsation to be an important force in affecting the flow of glymphatic drainage in the brain. In the meantime, AQP4 allows cytotoxic edema at the early phase of the cerebral ischemic stroke while removing vasogenic edema in the BBB at the late phase. Fine-tuning of AQP4 expression level in paravascular spaces is suggested to be used in the treatment of the brain edema caused by various stroke types.<sup>58</sup>

## Conclusion

In conclusion, this review article has attempted to gather the latest data on the brain glymphatic and immune systems. The function of the glymphatic system of the brain in the metabolite clearance has been broadly emphasized. Latest data leading to the diagnosis and treatment of the related diseases has been summarized. This paper has documented immunologic, molecular, physiologic, and anatomical relations of several neurotic pathologies including neurodegenerative diseases, tumors and related metastases, which will surely shed light on researchers dealing with this area.

## Conflict of Interest

The authors declare that they have no conflicts interest.

## References

1. Iliff JJ, Nedergaard M. Is there a cerebral lymphatic system? *Stroke*. 2013;44(Suppl. 1):93-5. doi:10.1161/STROKEAHA.112.678698
2. Iliff JJ, Goldman SA, Nedergaard M. Implications of the discovery of brain lymphatic pathways. *Lancet Neurol*. 2015;14(10):977-9. doi:10.1016/S1474-4422(15)00221-5
3. Mascagni P. Vasorum lymphaticorum corporis humani historia et ichnographia. Auctore Paulo Mascagni in Regio Senarum Lyceo publico anatomes professore. Senis: Ex typographia Pazzini Carli; 1787.
4. Retzius, Gustaf: Key A. Studien in der Anatomie des Nervensystems und das Bindegewebe - Digitised Book from the copyright-free holdings of the Bavarian State Library Munich 2007-2020 Image-based Similarity Search. 1875.
5. Sampson JH, Gunn MD, Fecci PE, Ashley DM. Brain immunology and immunotherapy in brain tumours. *Nat Rev Cancer*. 2020;20(1):12-25. doi:10.1038/s41568-019-0224-7
6. Cheng Y, Wang YJ. Meningeal Lymphatic Vessels: A Drain of the Brain Involved in Neurodegeneration? *Neurosci Bull*. 2020;36(5):557-60. doi:10.1007/s12264-019-00456-8
7. Sandrone S, Moreno-Zambrano D, Kipnis J, van Gijn J. A (delayed) history of the brain lymphatic system. *Nat Med*. 2019;25(4):538-40. doi:10.1038/s41591-019-0417-3
8. Louveau A, Smirnov I, Keyes TJ, Eccles JD, Rouhani SJ, Peske JD, et al. Structural and functional features of central nervous system lymphatic vessels. *Nature*. 2015;523(7560):337-41. doi:10.1038/nature14432
9. Mestre H, Mori Y, Nedergaard M. The Brain's Glymphatic System: Current Controversies. *Trends Neurosci*. 2020;43(7):458-66. doi:10.1016/j.tins.2020.04.003
10. Zamboni P. The discovery of the brain lymphatic system. *Veins Lymphat*. 2015;4 (2):5360. doi:10.4081/vl.2015.5360
11. Aspelund A, Antila S, Proulx ST, Karlsen TV, Karaman S, Detmar M, et al. A dural lymphatic vascular system that drains brain interstitial fluid and macromolecules. *J Exp Med*. 2015;212(7):991-9. doi:10.1084/jem.20142290
12. Ahn JH, Cho H, Kim JH, Kim SH, Ham JS, Park I, et al. Meningeal lymphatic vessels at the skull base drain cerebrospinal fluid. *Nature*. 2019;572(7767):62-6. doi:10.1038/s41586-019-1419-5
13. Gardner RC, Burke JF, Nettiksimmons J, Goldman S, Tanner CM, Yaffe K. Traumatic brain injury in later life increases risk for Parkinson disease. *Ann Neurol*. 2015;77(6):987-95. doi:10.1002/ana.24396
14. Kuo PH, Stuehm C, Squire S, Johnson K. Meningeal Lymphatic Vessel Flow Runs Countercurrent to Venous Flow in the Superior Sagittal Sinus of the Human Brain. *Tomography*. 2018;4(3):99-104. doi:10.18383/j.tom.2018.00013
15. De Angelis LC, Witte MH, Bellini T, Bernas M, Boccardo F, Ramenghi LA, et al. Brain lymphatic drainage system in fetus and newborn: Birth of a new era of exploration. *Lymphology*. 2018;51(4):140-7.
16. Gonzalez FF, Vexler ZS, Shimotake J, Ferriero DM. Cellular and molecular biology of hypoxic-ischemic encephalopathy. In: *Fetal and Neonatal Brain Injury*. Cambridge University Press; 2017. p. 36-49.
17. Teo JD, Morris MJ, Jones NM. Hypoxic postconditioning reduces microglial activation, astrocyte and caspase activity, and inflammatory markers after hypoxia-ischemia in the neonatal rat brain. *Pediatr Res*. 2015;77(6):757-64. doi:10.1038/pr.2015.47
18. Zhang X, Zhang X, Wang C, Li Y, Dong L, Cui L, et al. Neuroprotection of early and short-time applying berberine in the acute phase of cerebral ischemia: Up-regulated pAkt, pGSK and pCREB, down-regulated NF- $\kappa$ B expression, ameliorated BBB permeability. *Brain Res*. 2012;1459:61-70. doi:10.1016/j.brainres.2012.03.065
19. Li X, Bai R, Zhang J, Wang X. Effect of progesterone intervention on the dynamic changes of AQP-4 in hypoxic-ischaemic brain damage. *Int J Clin Exp Med*. 2015;8(10):18831-6.
20. Sun BL, Wang L hua, Yang T, Sun J yi, Mao L lei, Yang M feng, et al. Lymphatic drainage system of the brain: A novel target for intervention of neurological diseases. *Prog Neurobiol*. 2018;163:118-43. doi:10.1016/j.pneurobio.2017.08.007
21. Van Brussel I, Berneman ZN, Cools N. Optimizing dendritic cell-based immunotherapy: tackling the complexity of different arms of the immune system. *Mediators Inflamm*. 2012;2012:690643. doi:10.1155/2012/690643
22. Prins RM, Soto H, Konkankit V, Odesa SK, Eskin A, Yong WH, et al. Gene expression profile correlates with T-cell infiltration and relative survival in glioblastoma patients vaccinated with dendritic cell immunotherapy. *Clin Cancer Res*. 2011;17(6):1603-15. doi:10.1158/1078-0432.CCR-10-2563
23. Chertoff M, Shrivastava K, Gonzalez B, Acarin L, Gimenez-Llort L. Differential modulation of TREM2 protein during postnatal brain development in mice. *PLoS One*. 2013;8(8):e72083. doi:10.1371/journal.pone.0072083
24. Colonna M, Wang Y. TREM2 variants: New keys to decipher Alzheimer disease pathogenesis. *Nat Rev Neurosci*. 2016;17(4):201-7. doi:10.1038/nrn.2016.7
25. Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogaeva E,

- Majounie E, et al. TREM2 Variants in Alzheimer's Disease. *N Engl J Med.* 2013;368(2):117-27. doi:10.1056/NEJMoa1211851
26. Yaghmoor F. The Role of TREM2 in Alzheimer's Disease and Other Neurological Disorders. *J Alzheimer's Dis Park.* 2014;4(5):160. doi:10.4172/2161-0460.1000160
  27. Lim L, Sirichai P. Bone fractures: assessment and management. *Aust Dent J.* 2016;61:74-81. doi:10.1111/adj.12399
  28. Lyall K, Croen L, Daniels J, Fallin MD, Ladd-Acosta C, Lee BK, et al. The Changing Epidemiology of Autism Spectrum Disorders. *Annu Rev Public Health.* 2017;20:38:81-102. doi:10.1146/annurev-publhealth-031816-044318
  29. Lardenoije R, Iatrou A, Kenis G, Kompotis K, Steinbusch HW, Mastroeni D, et al. The epigenetics of aging and neurodegeneration. *Prog Neurobiol.* 2015;131:21-64. doi:10.1016/j.pneurobio.2015.05.002
  30. Bolte AC, Dutta AB, Hurt ME, Smirnov I, Kovacs MA, McKee CA, et al. Meningeal lymphatic dysfunction exacerbates traumatic brain injury pathogenesis. *Nat Commun.* 2020;11(1):4524. doi:10.1038/s41467-020-18113-4
  31. Louveau A, Da Mesquita S, Kipnis J. Lymphatics in neurological disorders: a neuro-lympho-vascular component of multiple sclerosis and Alzheimer's disease?. *Neuron.* 2016;91(5):957-73. doi:10.1016/j.neuron.2016.08.027
  32. Wang J, Jin WS, Bu X Le, Zeng F, Huang ZL, Li WW, et al. Physiological clearance of tau in the periphery and its therapeutic potential for tauopathies. *Acta Neuropathol.* 2018;136(4):525-36. doi:10.1007/s00401-018-1891-2
  33. Kim S, Kwon SH, Kam TI, Panicker N, Karuppagounder SS, Lee S, et al. Transneuronal Propagation of Pathologic  $\alpha$ -Synuclein from the Gut to the Brain Models Parkinson's Disease. *Neuron.* 2019;103(4):627-41. doi:10.1016/j.neuron.2019.05.035
  34. Smith AJ, Yao X, Dix JA, Jin BJ, Verkman AS. Test of the 'glymphatic' hypothesis demonstrates diffusive and aquaporin-4-independent solute transport in rodent brain parenchyma. *Elife.* 2017;6:e27679. doi:10.7554/eLife.27679
  35. Hablitz LM, Pla V, Giannetto M, Vinitsky HS, Staeger FF, Metcalfe T, et al. Circadian control of brain glymphatic and lymphatic fluid flow. *Nat Commun.* 2020;11(1):4411. doi:10.1038/s41467-020-18115-2
  36. Lundgaard I, Wang W, Eberhardt A, Vinitsky HS, Reeves BC, Peng S, et al. Beneficial effects of low alcohol exposure, but adverse effects of high alcohol intake on glymphatic function. *Sci Rep.* 2018;8(1):2246. doi:10.1038/s41598-018-20424-y
  37. Mendelsohn AR, Larrick JW. Sleep facilitates clearance of metabolites from the brain: glymphatic function in aging and neurodegenerative diseases. *Rejuvenation Res.* 2013;16(6):518-23. doi:10.1089/rej.2013.1530
  38. Xie L, Kang H, Xu Q, Chen MJ, Liao Y, Thiyagarajan M, et al. Sleep drives metabolite clearance from the adult brain. *Science.* 2013;342(6156):373-7. doi:10.1126/science.1241224
  39. Mosienko V, Teschemacher AG, Kasparov S. Is L-lactate a novel signaling molecule in the brain?. *J Cereb Blood Flow Metab.* 2015;35(7):1069-75. doi:10.1038/jcbfm.2015.77
  40. Scharf MT, Naidoo N, Zimmerman JE, Pack AI. The energy hypothesis of sleep revisited. *Prog Neurobiol.* 2008;86(3):264-80. doi:10.1016/j.pneurobio.2008.08.003
  41. Watts JC, Condello C, Stohr J, Oehler A, Lee J, DeArmond SJ, et al. Serial propagation of distinct strains of A $\beta$  prions from Alzheimer's disease patients. *Proc Natl Acad Sci U S A.* 2014;111(28):10323-8. doi:10.1073/pnas.1408900111
  42. Chen Z, Wilson MA. Deciphering neural codes of memory during sleep. *Trends Neurosci.* 2017;40(5):260-75. doi:10.1016/j.tins.2017.03.005
  43. Lee H, Xie L, Yu M, Kang H, Feng T, Deane R, et al. The effect of body posture on brain glymphatic transport. *J Neurosci.* 2015;35(31):11034-44. doi:10.1523/JNEUROSCI.1625-15.2015
  44. Andresen M, Hadi A, Petersen LG, Juhler M. Effect of postural changes on ICP in healthy and ill subjects. *Acta Neurochir.* 2014;157(1):109-13. doi:10.1007/s00701-014-2250-2
  45. O'Rourke DM, Nasrallah MP, Desai A, Melenhorst JJ, Mansfield K, Morrissette JJ, et al. A single dose of peripherally infused EGFRvIII-directed CAR T cells mediates antigen loss and induces adaptive resistance in patients with recurrent glioblastoma. *Sci Transl Med.* 2017;9(399):eaaa0984. doi:10.1126/scitranslmed.aaa0984
  46. Gupta A, Iadecola C. Impaired A $\beta$  clearance: a potential link between atherosclerosis and Alzheimer's disease. *Front Aging Neurosci.* 2015;7:115. doi:10.3389/fnagi.2015.00115
  47. Wang M, Ding F, Deng SY, Guo X, Wang W, Iliff JJ, et al. Focal solute trapping and global glymphatic pathway impairment in a murine model of multiple microinfarcts. *J Neurosci.* 2017;37(11):2870-7. doi:10.1523/JNEUROSCI.2112-16.2017
  48. Boluijt J, Meijers JC, Rinkel GJ, Vergouwen MD. Hemostasis and fibrinolysis in delayed cerebral ischemia after aneurysmal subarachnoid hemorrhage: a systematic review. *J Cereb Blood Flow Metab.* 2015;35(5):724-33. doi:10.1038/jcbfm.2015.13
  49. Benakis C, Garcia-Bonilla L, Iadecola C, Anrather J. The role of microglia and myeloid immune cells in acute cerebral ischemia. *Front Cell Neurosci.* 2015;8:461. doi:10.3389/fncel.2014.00461
  50. Fumagalli S, Perego C, Pischiutta F, Zanier ER, De Simoni MG. The ischemic environment drives microglia and macrophage function. *Front Neurol.* 2015;6:81. doi:10.3389/fneur.2015.00081
  51. Korkmaz A. The effect of helicobacter felis on macrophage polarization. *Istanbul Teknik Universitesi Fen Bilimleri Enstitusu Tez Calismasi.* 2015.
  52. Schilling M, Besselmann M, Leonhard C, Mueller M, Ringelstein EB, Kiefer R. Microglial activation precedes and predominates over macrophage infiltration in transient focal cerebral ischemia: a study in green fluorescent protein transgenic bone marrow chimeric mice. *Exp Neurol.* 2003;183(1):25-33. doi:10.1016/s0014-4886(03)00082-7
  53. Li M, Li Z, Ren H, Jin WN, Wood K, Liu Q, et al. Colony stimulating factor 1 receptor inhibition eliminates microglia and attenuates brain injury after intracerebral hemorrhage. *J Cereb Blood Flow Metab.* 2017;37(7):2383-95. doi:10.1177/0271678X16666551
  54. Sun BL, Xia ZL, Yan ZW, Chen YS, Yang MF. Effects of blockade of cerebral lymphatic drainage on cerebral ischemia after middle cerebral artery occlusion in rats. *Clin Hemorheol Microcirc.* 2000;23(2-4):321-5.
  55. Sun BL, Xia ZL, Wang JR, Yuan H, Li WX, Chen YS, et al. Effects of blockade of cerebral lymphatic drainage on regional cerebral blood flow and brain edema after subarachnoid hemorrhage. *Clin Hemorheol Microcirc.* 2006;34(1-2):227-32.
  56. Arbel-Ornath M, Hudry E, Eikermann-Haerter K, Hou S, Gregory JL, Zhao L, et al. Interstitial fluid drainage is impaired in ischemic stroke and Alzheimer's disease mouse models. *Acta neuropathologica.* 2013;126(3):353-64. doi:10.1007/s00401-013-1145-2
  57. Gaberel T, Gakuba C, Goulay R, De Lizarrondo SM, Hanouz JL, Emery E, et al. Impaired glymphatic perfusion after strokes revealed by contrast-enhanced MRI: A new target for fibrinolysis? *Stroke.* 2014;45(10):3092-6. doi:10.1161/STROKEAHA.114.006617
  58. Papadopoulos MC, Verkman AS. Aquaporin-4 gene disruption in mice reduces brain swelling and mortality in pneumococcal meningitis. *J Biol Chem.* 2005;280(14):13906-12. doi:10.1074/jbc.M413627200